Total number of printed pages – 6 MCA

Second Semester Examination - 2008

NUMERICAL METHODS

Full Marks – 70



Answer Question No. **1** which is compulsory and any **five** from the rest.

The figures in the right-hand margin indicate marks.

- 1. Answer the following questions : 2×10
 - (a) What is an error ? What are the different characteristics and types of error ?
 - (b) State the rules of rounding off decimal number correct upto n significant digits.
 - P.T.O.

- (c) Write -4.268106 and 0.00518789 in floating point form with 4 significant digits with round off.
- (d) State the basic difference between Secant method and method of false position in solving an equation.
- (e) What is the geometrical interpretation of Newton-Raphson method to solve an equation?
- (f) What is ill conditioning of a system of linear equations ? How can you overcome this problem ?
- (g) What is an eigen value of a matrix ? Why are eigen value problems important ?
- (h) Is Euler method to solve a differential equation accurate enough for practical problems? Can it be improved?
- What do you mean by single step and multi-step method to find the solution of a
- SCM 2006 2 Contd.





differential equation ? Give one example in each case.

- (j) How to calculate error in Simpson's 1/3 rule in evaluating an integral ?
- 2. (a) Determine a real root of the equation $x^{3.5} = 80$ using method of false position, correct upto three decimal places. 5
 - (b) What do you mean by the rate of convergence of an iterative method ? Determine the rate of convergence of Newton-Raphson method to find the solution of an equation.

5

 3. (a) Solve the following system of linear equations by Gauss elimination using partial pivoting: 5

$$2x + y + z = 1$$

$$5x + 2y + 2z = -4$$

$$3x + y + z = 5$$

SCM 2006 3 P.T.O.

(b) Solve the following system of linear equations by Gauss-Seidel method correct upto two decimal places : 5

10x + 2y - z = 27-3x - 6y + 2z = -61.5

$$x + y + 5z = -21.5$$

4. (a) Using least square regression, find a straight line to the following given data: 5

Х	:	1	2	3	4	5	6	7	8	9
v		1	15	2	3	Δ	5	8	10	13

- (b) Using centered difference approximation, estimate the first and second order derivative of $y = e^x$ at x = 2 for the step size h = 0.1. 5
- 5. (a) Derive and estimate the error of Trapezoidal rule for numerical integration of a function f in the range [a, b].
 5
 SCM 2006
 4
 Contd.

(b) Using Simpson's 3/8 rule evaluate the following integral by using 9 sub-intervals of equal width: 5

$$\sum_{0}^{\frac{\lambda}{2}} \mathbf{Z}_{\text{in}} x \, dx.$$

6. (a) Solve the following differential equation by using modified Euler's method for y(4.1)and y(4.2), taking h = 01 5

$$5x\frac{dy}{dx}+y^2-2=0.$$

(b) Solve the following differential equation for y(0.1) and y(0.2) using Runge-Kutta method of fourth order.

$$\frac{dy}{dx} = \frac{1}{x+y} \text{ where } y(0) = 1.$$

7. (a) Find all eigen values and eigen vectors of the matrix 5

5

SCM 2006

(b) Find the inverse of the following matrix by using LU decomposition.5

- 8. (a) Evaluate the polynomial $y = x^3 7x^2 + 8x 0.35$ at x = 1.37. Use 3-digit arithmetic with chopping. Also evaluate the percent relative error. 5
 - (b) Discuss about stability and condition of a mathematical problem. Also compute and interpret the condition number for

$$f(x) = \tan x$$
 for $x = \frac{\pi}{2} + 0.1 \frac{\pi}{2}$. 5

SCM 2006

6

– C